

Review on vocational predictors: a systematic review of predictors of vocational outcomes among individuals with schizophrenia: an update since 1998

Hector W.H. Tsang, Ada Y. Leung, Raymond C.K. Chung, Morris Bell, Wai-Ming Cheung

Objective: Predictors of employment outcomes of individuals with schizophrenia have continued to be studied over the past decade with implications for the development of vocational interventions to help the mentally ill get and keep jobs.

Methods: A total of 62 relevant studies since 1998 were systematically reviewed by means of meta-analysis and frequency counts. Frequency count allowed all 62 studies to be included, whereas the meta-analysis excluded studies with inadequate information but made it possible to estimate the magnitude of effects.

Results: Both methods resulted in similar findings. In contrast to an earlier review, cognitive functioning received overwhelming support as a significant predictor. Other significant predictors included education, negative symptoms, social support and skills, age, work history (previous history of successful employment), and rehabilitation service to restore community functioning and well-being by occupational therapists, psychiatrists, psychologists, social workers and other mental health professionals. Positive symptoms, substance abuse, gender and hospitalization history were found to be non-significant predictors. The frequency count did not support marital status as a significant predictor but the meta-analysis did.

Conclusions: This review highlights increasing sophistication in understanding the links between individual characteristics and functional impairments. It also suggests that more research is needed into other potentially important predictors that may be changeable and relate to recovery. These include attitudes and beliefs about disability payments and psychological processes such as self-stigmatization, negative beliefs, and social skills deficits for which intervention may be possible.

Key words: employment outcome, frequency, meta-analysis, rehabilitation, schizophrenia.

Australian and New Zealand Journal of Psychiatry 2010; 44:495–504

Hector W.H. Tsang, Associate Professor (Correspondence); Ada Y. Leung, Project Associate; Raymond C.K. Chung, Scientific Officer

Neuropsychiatric Rehabilitation Laboratory, Department of Rehabilitation Sciences, The Hong Kong Polytechnic University, Hung Hom, Hong Kong, China. Email: rshtsang@inet.polyu.edu.hk

Morris Bell, Professor

Rehabilitation Research & Development Service, US Department of Veterans Affairs & Department of Psychiatry, Yale University School of Medicine, New Haven, Connecticut, USA

Wai-Ming Cheung, Assistant Professor

Faculty of Education, University of Hong Kong, Hong Kong, China

Received 8 October 2009; accepted 16 March 2010.

When Anthony and Jansen [1] first reported that psychiatric symptomatology and diagnosis were poor predictors of future work performance for the mentally ill, it was inspiring to many in psychiatric rehabilitation. Their findings suggested that the vocational potential of mentally ill individuals should not be prejudged based on these factors. Indeed, it appeared that prior employment history was the best and most reliable predictor of future vocational performance along with work adjustment skills, ability to 'get along' or function socially with others, and ego strengths or self-concept. The Anthony

and Jansen [1] review article had a significant impact on the development of psychiatric rehabilitation. First, it highlighted the importance of functional outcomes among people with schizophrenia; and, second, its findings had direct effects on the development of vocational interventions to help the mentally ill get and keep their jobs.

In 1984 there was little else that Anthony and Jansen [1] could say about what features of psychiatric illness might be relevant to rehabilitation because the discovery of how mental illness affects vocational functioning had to await developments in the study of psychopathology and functional impairment. Research along these lines continued and was enriched by more sophisticated models of the relationship between psychopathology and function that importantly included the distinction between positive and negative symptoms and the role of neurocognitive and social cognitive impairments. For instance, Bell, Tsang and colleagues [2] studied the relationship between cognitive impairments and rate of symptom improvement in a 26-week work therapy programme from 1998 to 2004. They found that logical memory was a significant predictor of symptom change and work performance. This implied that improvements in cognitive functioning might have an impact on both symptoms and work functioning.

In view of the rapidly growing body of knowledge, we published a review paper that summarized the consistent predictors of vocational performance of psychiatric patients based on 35 relevant studies from 1985 to 1997 [3]. Our review concluded that premorbid occupational functioning, work history, and social skills were consistent predictors. Symptomatology and diagnosis continued to be inconclusive as to their predictive power. Cognitive functioning and family relationships, although largely neglected by researchers, were found to be significant predictors. Since its release in 2000, this article has been frequently cited, in part because vocational rehabilitation has become a legitimate focus for community mental health centres serving individuals with severe mental illnesses [4–7]. A study on perceptions among psychiatric practitioners on remission and recovery in schizophrenia [8] found that more than half (62%) of the respondents endorsed ‘returning to work’ as the core element of recovery. Another study [9] reported that full recovery can only be achieved when a person holds a steady job. More importantly, studies have shown that most individuals with severe mental illness express a strong desire to work [10,11].

The study of predictors of vocational functioning in schizophrenia has also been important as clues to underlying mechanisms of illness that may explain social dysfunction more generally. In addition, the discovery of the predictive power of neurocognitive [12–14,15–18] social

cognitive [19] and social skills [12,20–22] deficits has led to the development of interventions that target these deficits in the interest of improving vocational outcomes.

In line with the increasing interest in helping clients improve their employability, the volume of studies that examine predictors of vocational outcomes has continued to expand. There are two characteristic features of recent publications pertaining to vocational outcomes. The first is that cognitive function has received more and more attention. This may be due to the seminal article by Green (1996) [23] which linked neurocognitive abilities to functional outcome for people with schizophrenia which was subsequently supported by empirical studies [24–26]. Another change we observed in the more recent literature is an increase in the sophistication of statistical modelling such as path analysis (PA) and structural equation modelling (SEM) to reveal the direct and indirect contributions of multiple variables in relation to vocational outcomes [27–29]. Despite the availability of this body of new data, there have not been many comprehensive reviews on predictors of vocational outcomes since our own publication in 2000. From 2000, our search revealed six systematic reviews [30–35] on vocational rehabilitation and severe mental illness. However, all of these reviews have limitations and did not give a comprehensive account of vocational predictors for people with schizophrenia based on available information. Our present review thus fills the gap in the literature as we cover studies which were published from 1998 to 2008. Although meta-analysis has been widely used to pool effect sizes of Randomized Clinical Trials (RCTs) and come up with a relevant statistic which summarizes the effectiveness of a particular clinical intervention [36–43], there is to date no attempt to apply similar methods to the study of predictors. In this paper we employed the meta-analytical method using the generic inverse variance [44].

Method

Data searching and initial screening criteria

To collect literature pertaining to the significant predictors of vocational outcomes, a comprehensive electronic search was performed over the period from January 1998 to 2008 with computerized electronic databases (EBSCOhost, InterScience, Journals@Ovid, MEDLINE, MD Consult, PsycINFO, ScienceDirect, Science Citation Index/Social Science Citation Index), using the following keywords ‘schizophrenia’ or ‘mental illness’ or ‘psychiatric disability’ or ‘psychiatric disorder’ and ‘employment’ or ‘vocational outcome’ and ‘prediction’ or ‘predictors’. Studies were further filtered by visual inspection to avoid duplication of articles across different databases. A total of 439 articles (260 from MEDLINE, 120 from Science Citation Index and

Social Science Citation Index, 10 from EBSCOhost, 6 from Science Direct, 7 from Journals@Ovid, 18 from MD Consult, 16 from PsycINFO, and 2 from Interscience) were extracted. A study was included in our meta-analysis if: (i) the majority (more than 50%) of the participants were diagnosed as suffering from schizophrenia by certified psychiatrists or by DSM-III, DSM-III-R, or DSM-IV criteria; (ii) the theme was mainly on identification of predictors of vocational outcome; and (iii) the method was empirical and quantitative in nature. The study was excluded if it was a qualitative study. Following the above screening criteria, 72 articles (31 from MEDLINE, 22 from Science Citation Index and Social Science Citation Index, 9 from PsycINFO, 6 from EBSCOhost, 3 from MD Consult, and 1 from Science Direct) were selected for a detailed review at the second stage.

Second stage screening

At the second stage screening, two qualified reviewers with a Master's degree in clinical psychology or occupational therapy screened through all eligible studies and decided whether each of them should be included for data synthesis in the next stage based on the criteria below. As to methodology, at least one of the objectives was to identify significant predictors of vocational outcomes. The criteria for subject recruitment were clearly defined and the diagnosis of schizophrenia was valid. The assessments for employment status and correlation/predictor variables were validated. Finally, the assessment was performed by independent and qualified assessors. Regarding the criteria for data analysis, only studies using multivariate analysis such as logistic regression, multiple regression, MANOVA, path analysis, structural equation model, etc., on derivation of predictors were included [45]. In addition, we included studies which involved at least five subjects per predictor variable [46], had less than 20% attrition rate [47], and had a priori power analysis on sample size determination [48]. On the other hand, we excluded studies using proportional hazards models with hazard rate estimates quoted. The reason was that method for conversion of hazard rate to odds ratio (OR) was unavailable. If the same research group published more than one paper for the same purpose with the same population, only the most recently published article was included for review [46]. If the same research group published more than one paper with the same population but different purposes, all the papers were included.

After screening by the two raters, 53 articles were selected for extensive review and data synthesis. The concordance rate [49] between the two raters was 80%. After extracting the 53 articles, we found that some papers were missed because they might not be found by the keywords in the databases. Therefore, we supplemented the 53 articles with an additional 9 articles which are also relevant to our current review on predictors of vocational outcome for people with schizophrenia. A similar method was used in McGurk and Mueser's review article [33]. Eventually 62 articles were included in the systematic review. Figure 1 summarizes the screening process leading to the 62 studies for a detailed meta-analysis.

Data analysis

We used two methods for data analysis. First, we replicated the method we used in our earlier review [3] by frequency counts to summarize studies that reported whether an individual predictor was a significant predictor or not. Second, we conducted a meta-analysis on the 62 articles. As to meta-analysis, we aimed to transform the results from

the 62 studies into a common metric so that statistical comparisons could be made across the predictors and outcomes [50]. Most studies that we reviewed used regression analysis for either continuous or categorical predictors. In studies involving continuous predictors, standardized regression weights (beta coefficients) were reported. In studies using categorical predictors, the most common effect size was reported as odds ratio. Based on Portney and Watkins [51], odds ratio is a measure of effect size defined as the ratio of the odds of an event occurring in one group to the odds of it occurring in another group. An odds ratio of 1 indicates that the condition or event under study is equally likely in both groups. An odds ratio greater than 1 indicates that the condition or event is more likely in the first group. On the contrary, an odds ratio less than 1 indicates that the condition or event is less likely in the first group. In this meta-analysis, we used odds ratios as the effect estimates for each study because of the ease of interpretation in applied settings and strong mathematical properties [52]. The odds ratio between each predictor variable and each vocational outcome for each study was estimated by cell counts in a 2×2 contingency table [53]. In some studies they were extracted directly from the articles without further computation. If odds ratios were not reported in a study, the following three methods were employed to transform the effect size information into a common odds ratio metric, making the results of the studies comparable across different analytical methods [50]. First, we transformed the correlation coefficient into odds ratio using the formula suggested by Digby (1983) [54] if the correlation coefficient estimates were given instead. Second, regression coefficients of predictor variables were converted into Pearson's correlation coefficients. If the results were based on logistic regression models, the regression coefficient of each predictor was converted into odds ratio by exponential transformation. Finally, 41 studies listing only the p-value of each predictor variable without mentioning the magnitudes for any type of effect or not providing results convertible to odds ratio were excluded from the meta-analysis.

Standard errors of the effect size estimates were computed either by conversion of the test statistic or by confidence interval. The test statistic was defined as the natural log odds ratio divided by its standard error [53] and was converted into p-values [51]. By proper transformations, the standard error of natural log odds ratios was obtained. Another way of estimating standard error of the effect estimate was utilization of the width of the individual 95% confidence interval [55]. In addition, the definition of the odds ratio across studies was standardized for directionality. Following these conversions, meta-analytic techniques were used to estimate the overall effect sizes of odds ratio [56]. Regarding the meta-analysis of each predictor on each vocational variable for all eligible studies, the generic inverse variance approach [57] was used to pool all individual effect size estimates for computation of the overall effect size estimates by assuming fixed-effect models with the use of Review Manager 4.2.9 (The Cochrane Collaboration, 2005) [55]. For study with two experimental groups [58], weighted average of correlations was used to obtain a single effect size. Along with this, studies with multiple predictor variables in a single conceptual domain, average across all the variables was computed to obtain the single effect size [59–63]. The pooled effect size estimate of each predictor on each vocational outcome was regarded as statistically significant if its 95% confidence interval did not cover the value of one. Furthermore, cognitive functioning was divided into three aspects (i.e. executive functioning and general intelligence, attention and working memory, and verbal and visual memory) so that cognitive tests could be pulled together for meta-analysis. Odds ratios of the scores of Wisconsin Card




Figure 1. Flowchart of Review Articles Selection Process.

Sorting Test (WCST) and Wechsler Adult Intelligence Scale (WAIS) in predicting vocational outcomes were grouped for determining the effect size of executive functioning and general intelligence. Odds ratios of Trail Making Tests, Digit Span, and Repeatable Battery for the Assessment of Neuropsychological Status (RBANS)'s attention index were pooled for attention and working memory. Finally, the California Verbal Learning Test (CVLT), RBAN's immediate memory index, and RBAN's delayed memory index were pooled for verbal and visual memory.

Results

Categorization of predictors

The significant predictors identified from the 62 studies were grouped into 15 categories by the first and second authors (H.W.T.S. and A.Y.L.),

with 100% concordance. The 15 categories were further broken down into 41 individual predictors. They were psychiatric symptomatology (positive symptom, negative symptom, general psychopathology), demographic characteristic (age, gender, education, ethnicity, marital status, socio-economic status), cognitive functioning, functional history and attitudes (premorbid functioning, attitude towards work, work adjustment skills, work history), substance abuse, public support/disability income (i.e. publicly funded disability support scheme such as a pension, etc.), rehabilitation services to restore community functioning and well-being by occupational therapists, psychiatrists, psychologists, social workers and mental health professionals, social features (social support and skills, psychosocial stressor), work income, family history, hospitalization history, emotion, intra-psyche functioning, medical features (medical condition or disability which refers to impaired body functions requiring medical attention, diagnosis, atypical antipsychotic medication, first generation antipsychotic medication, medication adher-

ence, length of illness, duration of untreated psychosis, age of onset of illness, course of illness, remission) and others (duration of military service before admission, work quality, work habit, family functioning, family social economic status, therapeutic relationship, self-esteem).

Description of studies

Table 1 shows the number of studies that cited each of the predictors as significant or non-significant. Results show that cognitive functioning was the most intensively studied variable and received overwhelming support (N=15) as a significant predictor. However, at the same time, seven studies reported it as a non-significant predictor. The next two variables that received much attention from the studies and were mostly cited as significant predictors included education (N=12) and negative symptoms (N=10). But these two predictors were simultaneously regarded as non-significant by seven and nine studies respectively. Social support and skills (N=8), medical condition or disability (N=8), work history (N=7) and rehabilitation service (N=7) were also reported as significant predictors. Five studies unanimously reported public support or disability income as a significant negative predictor. Twelve predictors (socio-economic status, premorbid functioning, work adjustment skills, family history, emotion, intra-psychic functioning, atypical antipsychotic medication, medication non-adherence, length of illness, duration of untreated psychosis, duration of military service before admission, and therapeutic relationship) received support from only one study as being significant.

Positive symptoms (N=11), diagnosis (N=7) and substance abuse (N=6) received the most support as non-significant predictors; but positive symptoms were regarded as significant predictors by five studies as well. Similarly, both gender and hospitalization had support from nine studies as non-significant and five as significant. Furthermore, ethnicity, marital status, length of illness, and age of onset all received support as non-significant predictors by five studies.

The most contradictory results were found in negative symptoms and age, they received similar support as significant (N=10 for negative symptom, and N=7 for age) and non-significant predictor (N=9 for negative symptom, and N=8 for age).

Meta-analysis

Table 2 shows the number of studies used and the results of the pooled effect size of the predictors. The effect sizes of studies with continuous predictors and categorical predictors are pooled separately. There were fewer studies that had the information required for inclusion in this form of analysis: 21 studies were included in the meta-analysis. Negative symptoms of age, education, marital status, work history, public support, diagnosis, and executive functioning and general intelligence were all significant predictors with the highest OR for marital status and diagnosis. Positive symptoms of gender, ethnicity, hospitalization history, attention and working memory, and verbal and visual memory were non-significant predictors.

Discussion

Our review of published research since 1998 has revealed continued refinement and development in the

investigation of predictors. In particular there has been a surge of interest in the role of cognitive function. In contrast to our earlier report, results show that cognitive functioning was the most intensively studied variable, and it received overwhelming support (N=15) as a significant predictor. Although all the predictors in the current review had general representation in the earlier literature, there was greater specificity, particularly in regard to distinguishing between negative symptoms and positive symptoms. Social factors that might be barriers to employment did not receive much attention, but five studies unanimously reported public support or disability income as a significant negative predictor. This reflects that people with mental illness who are on a publicly funded disability support scheme such as a pension scheme may predict poorer employment outcome. Psychological factors such as self-stigmatizing beliefs about their own mental illness, negative cognitions about themselves as workers and the absence of positive psychological factors [64] such as hopefulness and optimism are currently receiving some attention as predictors of social functioning in the context of the recovery movement, but they have yet to find their way into formal studies of predictors of vocational outcome.

This report provided two approaches to reviewing the literature. The frequency count of studies allowed us to include more studies and variables than was the case for the meta-analysis. However, the frequency count method only showed the number of studies of the significant and non-significant predictors instead of counting the number of subjects in a study corresponding to a predictor. The meta-analysis had the advantage of providing odds ratios, which made comparisons possible among variables; for example, marital status emerged as having the highest odds ratio, whereas work history had a significant but much smaller effect. We used both methods so that the readers may cross-check the results of each method. However, as both methods have advantages and limitations, cautions have to be made accordingly.

The meta-analysis for both categorical and continuous variables reported that cognitive functioning (specifically executive function and general intelligence), negative symptom, and age were significant predictors, which agreed with the frequency counts. Similarly, work history, education, and public support/disability income were also found to be statistically significant as categorical variables. Gender and hospitalization history were found to be non-significant predictors, which paralleled the result of the frequency count. Marital status was an exception to the consistency in findings between frequency counts and meta-analysis. Based on meta-analysis, marital status was found to be a significant predictor but the frequency count showed one study supporting it

Table 1. Frequency counts of studies on significant or non-significant predictors

Predictors	Significant predictors as indicated by the number of studies reviewed	Non-significant predictors as indicated by the number of studies reviewed
	N	N
Psychiatric symptomatology		
Positive symptom	5	11
Negative symptom	10	9
General psychopathology	2	1
Demographic characteristics		
Age	7	8
Gender	5	9
Education	12	7
Ethnicity	2	5
Marital status	2	5
Socioeconomic status	1	0
Cognitive functioning	15	7
Functional history and attitudes		
Premorbid functioning	1	1
Attitude towards work	2	0
Work adjustment skills	1	1
Work history	7	1
Substance abuse	2	6
Public support/disability income	5	0
Rehabilitation services	7	2
Social features		
Social support and skills	8	3
Psychosocial stressor	2	0
Work income	0	1
Family history	1	2
Hospitalization history	5	9
Emotion	1	0
Intrapsychic functioning	1	0
Medical features		
Medical condition or disability	8	4
Diagnosis	3	7
Atypical antipsychotic medication	1	3
First generation antipsychotic medication	0	3
Medication non-adherence	1	1
Length of illness	1	5
Duration of untreated psychosis	1	0
Age of onset of illness	3	5
Course of illness	3	0
Remission	1	1
Others		
Duration of military service before admission	1	0
Work quality	2	0
Work habit	2	1
Family functioning	0	1
Family social economic status	0	1
Therapeutic relationship	1	0
Self-esteem	0	1

and three studies not supporting it. This contradiction may likely have been because only two of the four studies used in the frequency count could be included in the meta-analysis. Also, the meta-analysis could weigh and

combine findings so that the strongly positive finding in one study, when combined with the weak negative finding in the other, still yielded a positive overall effect. Had all four studies been suitable for inclusion, the meta-analytic

approach would have been superior to the frequency count in fairly evaluating the predictive value of marital status.

In this review we wanted to weigh the scientific evidence regarding individual predictors because they have potential significance for developing vocational interventions. Some variables such as education may reflect pre-morbid characteristics such as social class, educational opportunity and the age at which psychiatric illness may have begun to impair social functioning (which may well precede onset of illness). These variables are aspects of the person's past and are not changeable. They are relevant to rehabilitation primarily in terms of appropriate accommodations. Some variables, such as negative symptoms and cognitive functioning represent features of illness that directly affect current functioning and may be changeable. They deserve the attention they are beginning to get as targets for novel medications and remediation strategies. Vocational programs may wish to consider integrating these treatments into their rehabilitation plans in a coordinated effort to improve function through the synergy of new treatments and work opportunity. While medications targeting negative symptoms and cognitive enhancement have yet to be approved, a few experimental studies using cognitive remediation with

vocational rehabilitation have been reported with good results [18,65,66]. Such integration of treatment with rehabilitation is consistent with the individual placement and support (IPS) model [67,68], which recognizes the important role that a team approach between clinicians and vocational specialists can play in successful vocational outcomes.

This review also shows that more research is needed into other potentially important predictors that may be changeable and relate to recovery. These include social cognition, social policy, and attitudes related to disability income, and psychological processes such as self-stigmatization, negative beliefs, and social skills deficits for which intervention may be possible [8,20,26,69]. This is particularly true for social cognition which has received more and more attention among researchers because of its relevance to vocational outcomes. For instance, poorer social cognition could lead to social discomfort on the job which in turn could lead to poorer work rehabilitation outcomes [26,70]. By understanding the relationships between individual characteristics and vocational outcomes, the field can gain valuable direction for the development and refinement of interventions that help people overcome their personal barriers to vocational success.

Table 2. Results of pooled analysis of outcome parameters

Outcome comparisons		Studies	Participants	Statistical method	
				OR#	95% CI
Categorical outcomes (competitive employment)					
Psychiatric symptomatology	Negative symptom	3[71–73]	2664	0.91	(0.88, 0.94)*
	Positive symptom	3[72–74]	4757	1.01	(0.98, 1.03)
Demographic characteristic	Younger age	8[60,61,71,73,75–78]	5788	0.78	(0.75, 0.81)*
	Female	4[60,78–80]	1531	0.75	(0.51, 1.10)
	Education	9[6,60,61,63,71,73,74,78,79]	6367	0.94	(0.89, 0.99)*
	Marital status (Married/cohabiting)	3[78–80]	800	1.96	(1.85, 2.08)*
Functional features	Ethnicity	2[75,78]	1422	0.86	(0.39, 1.91)
	Work history	5[71,75,78,81,82]	2148	1.06	(1.02, 1.11)*
Public support/disability income		2[6,81]	1931	0.29	(0.26, 0.32)*
Hospitalization history		4[62,63,73,77]	3817	1.05	(1.0, 1.11)
Medical features	Diagnosis	2[61,82]	1055	1.6	(1.15, 2.23)*
Continuous outcome (hours worked)					
Psychiatric symptomatology	Negative symptom	5[58,62,73,83,84]	3883	0.89	(0.87, 0.91)*
Demographic characteristic	Age	2[75,83]	1465	1.03	(1.01, 1.05)*
Gender	Male	2[73,75]	3599	1.13	(1.07, 1.20)*
Cognitive functioning	Executive and intelligence	4[58,59,72,84]	270	0.59	(0.40, 0.89)*
	Attention and working memory	3[58,59,84]	182	1.36	(0.61, 3.04)
	Verbal and visual memory	2[59,84]	70	1.28	(0.42, 3.85)

#Odds ratio; *Significant predictor.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

1. Anthony WA, Jansen MA. Predicting the vocational capacity of the chronically mentally ill. *Am Psychol* 1984; 39:537–544.
2. Bell MD, Tsang HW, Greig T, Bryson G. Cognitive predictors of symptom change for participants in vocational rehabilitation. *Schizophr Res* 2007; 96:162–168.
3. Tsang HWH, Lam P, Ng B, Leung, O. Predictors of employment 3. outcome of people with psychiatric disabilities: a review of the literature since mid 80s. *J Rehabil* 2000; 66:19–31.
4. Bond GR, Salyers MP, Dincin J, Drake R, Becker DR, Fraser VV, Haines M. A randomized controlled trial comparing two vocational models for persons with severe mental illness. *J Consult Clin Psychol* 2007; 75:968–982.
5. Drake RE, Becker DR, Bond GR. Recent research on vocational rehabilitation for persons with severe mental illness. *Curr Opin Psychiatry* 2003; 16:451–455.
6. Rosenheck R, Leslie D, Keefe R, McEvoy J, Swartz M, Perkins D, Stroup S, Hsiao JK, Lieberman J. Barriers to employment for people with schizophrenia. *Am J Psychiatry* 2006; 163: 411–417.
7. Rosenthal DA, Dalton JA, Gervy R. Analyzing vocational outcomes of individuals with psychiatric disabilities who received state vocational rehabilitation services: a data mining approach. *Int J Soc Psychiatry* 2007; 53:357–368.
8. Tsang HW, Chen EY. Perceptions on remission and recovery in schizophrenia. *Psychopathology* 2007; 40:469.
9. Ng RM, Pearson V, Lam M, Law CW, Chiu CP, Chen EY. What does recovery from schizophrenia mean? Perceptions of long-term patients. *Int J Soc Psychiatry* 2008; 54:118–130.
10. McQuilken M, Zahniser JH, Novak J, Starks RD, Olmos A, Bond GR. The Work Project Survey: consumer perspectives on work. *J Vocational Rehabil* 2003; 18:59–68.
11. McGurk SR, Mueser KT, Pascaris A. Cognitive training and supported employment for persons with severe mental illness: one-year results from a randomized controlled trial. *Schizophr Bull* 2005; 31:898–909.
12. Bell MD, Choi J, Lysaker P. Psychological interventions to improve work outcomes for people with psychiatric disabilities. *J Norwegian Psychol Assoc* 2007; 44:606–617.
13. Bell MD, Bryson G. Work rehabilitation in schizophrenia: does cognitive impairment limit improvement? *Schizophr Bull* 2001; 27:269–279.
14. Bell M, Fiszdon J, Greig T, Wexler B, Bryson G. Neurocognitive enhancement therapy with work therapy in schizophrenia: six-month follow-up of neuropsychological performance. *J Rehabil Res Dev* 2007; 44:761–770.
15. Bell MD, Zito W, Greig T, Wexler BE. Neurocognitive enhancement therapy with vocational services: work outcomes at two-year follow-up. *Schizophr Res* 2008; 105:18–29.
16. Wexler BE, Bell MD. Cognitive remediation and vocational rehabilitation for schizophrenia. *Schizophr Bull* 2005; 31: 931–941.
17. Bell MD, Bryson GJ, Greig TC, Fiszdon JM, Wexler BE. Neurocognitive enhancement therapy with work therapy: Productivity outcomes at 6- and 12-month follow-ups. *J Rehabil Res Dev* 2005; 42:829–838.
18. Greig TC, Zito W, Wexler BE, Fiszdon J, Bell MD. Improved cognitive function in schizophrenia after one year of cognitive training and vocational services. *Schizophr Res* 2007; 96:156–161.
19. Bell MD, Zito W, Greig T, Wexler B. Neurocognitive enhancement therapy and competitive employment in schizophrenia: effects on clients with poor community functioning. *Am Journal Psychiatr Rehabil* 2008; 11:109–122.
20. Tsang HWH. Augmenting vocational outcomes of supported employment by social skills training. *J Rehabil* 2003; 69: 25–30.
21. Tsang HWH. Enhancing employment opportunities of people with mental illness through integrated supported employment: *Health Services Research Committee Report*, 2007: 1–6.
22. Tsang HWH, Chan A, Wong A, Liberman RP. Vocational outcomes of an integrated supported employment program for individuals with persistent and severe mental illness. *J Behav Ther Exp Psychiatry* 2009; 40:292–305.
23. Green MS. What are the functional consequences of neurocognitive deficits in schizophrenia. *AM J Psychiatry* 1996; 153:321–330.
24. Bell M, Bryson G, Greig T, Corcoran C, Wexler BE. Neurocognitive enhancement therapy with work therapy: effects on neuropsychological test performance. *Arch Gen Psychiatry* 2001; 58:763–768.
25. Bryson G, Bell MD. Initial and final work performance in schizophrenia: cognitive and symptom predictors. *J Nerv Ment Dis* 2003; 191:87–92.
26. Bell M, Tsang HW, Greig TC, Bryson GJ. Neurocognition, social cognition, perceived social discomfort, and vocational outcomes in schizophrenia. *Schizophr Bull* 2009; 35:738–747.
27. Wu HC. Predicting subjective quality of life in workers with severe psychiatric disabilities. *Community Ment Health J* 2008; 44:135–146.
28. Brekke J, Kay DD, Lee KS, Green MF. Biosocial pathways to functional outcome in schizophrenia. *Schizophr Res* 2005; 80:213–225.
29. Bellack AS, Green MF, Cook JA, et al. Assessment of community functioning in people with schizophrenia and other severe mental illnesses: a white paper based on an NIMH-sponsored workshop. *Schizophr Bull* 2007; 33:805–822.
30. Wewiorski NJ, Fabian ES. Association between demographic and diagnostic factors and employment outcomes for people with psychiatric disabilities: a synthesis of recent research. *Ment Health Serv Res* 2004; 6:9–21.
31. Cook JA, Razzano L. Vocational rehabilitation for persons with schizophrenia: recent research and implications for practice. *Schizophr Bull* 2000; 26:87–103.
32. Christensen TO. The influence of neurocognitive dysfunctions on work capacity in schizophrenia patients: a systematic review of the literature. *Int J Psychiatry in Clin Pract* 2007; 11:89–101.
33. McGurk SR, Mueser KT. Cognitive functioning, symptoms, and work in supported employment: a review and heuristic model. *Schizophr Res* 2004; 70:147–173.
34. Bond GR, Drake RE. Predictors of competitive employment among patients with schizophrenia. *Curr Opin Psychiatry* 2008; 21:362–369.
35. Michon HW, van Weeghel J, Kroon H, Schene AH. Person-related predictors of employment outcomes after participation in psychiatric vocational rehabilitation programmes: a systematic review. *Soc Psychiatry Psychiatr Epidemiol* 2005; 40:408–416.
36. Davis JM, Chen N, Glick ID. A meta-analysis of the efficacy of second-generation antipsychotics. *Arch Gen Psychiatry* 2003; 60:553–64.

37. Morris SB, DeShon RP. Combining effect size estimates in meta-analysis with repeated measures and independent-groups designs. *Psychol Methods* 2002; 7:105–125.
38. Saint S, Veenstra DL, Sullivan SD. The use of meta-analysis in cost-effectiveness analysis. *Pharmacoeconomics* 1999; 15:1–8.
39. Takkouche B, Cadarso-Suarez C, Spiegelman D. Evaluation of old and new tests of heterogeneity in epidemiologic meta-analysis. *Am J Epidemiol* 1999; 150:206–215.
40. Leucht S, Wahlbeck K, Hamann J, Kissling W. New generation antipsychotics versus low-potency conventional antipsychotics: a systematic review and meta-analysis. *Lancet* 2003; 361:1581–9.
41. Tuunainen A, Wahlbeck K, Gilbody S. Newer atypical antipsychotic medication in comparison to clozapine: a systematic review of randomized trials. *Schizophr Res* 2002; 56:1–10.
42. Rector NA, Beck AT. Cognitive behavioral therapy for schizophrenia: an empirical review. *J Nerv Ment Dis* 2001; 189:278–87.
43. Zimmermann G, Favrod J, Trieu VH, Pomini V. The effect of cognitive behavioral treatment on the positive symptoms of schizophrenia spectrum disorders: a meta-analysis. *Schizophr Res* 2005; 77:1–9.
44. Bonett DG. Transforming odds ratios into correlations for meta-analytic research. *Am Psychol* 2007; 62:254–255.
45. Adams E, Karen F, Elaine A. A systematic review of clinical predictors of outcomes in adults with recent major limb amputation. *VA OPCS Technol Assessment Progam. USA: United States Department of Veterans Affairs, 2005.*
46. Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 1996; 49:1373–1379.
47. Pincus T, Burton AK, Vogel S, Field AP. A systematic review of psychological factors as predictors of chronicity/disability in prospective cohorts of low back pain. *Spine* 2002; pp. E109–E120.
48. Lawrence R, Bradshaw T, Mairs H. Group cognitive behavioural therapy for schizophrenia: a systematic review of the literature. *J Psychiatr Ment Health Nurs* 2006; 13:673–681.
49. Edwin JCG. Concordance Rate. *Encyclopedia of Statistics in Behavioral Science* 2005; 1:359.
50. Roberts BW, Kuncel NR, Shiner R, Caspi A, Goldberg LR. The power of personality: the comparative validity of personality traits, socioeconomic status, and cognitive ability for predicting important life outcomes. *Perspectives Psychol Sci* 2007; 2:313–344.
51. Portney LG, Watkins MP. *Foundation of clinical research: applications to practice.* Upper Saddle River (NJ): Prentice Hall, 2000.
52. Mantel N, Brown C, Byar DP. Tests for homogeneity of effect in an epidemiologic investigation. *Am J Epidemiol* 1977; 106:125–129.
53. Altman DG. *Practical Statistics for Medical research.* Florida: Boca Roton, 1991.
54. Digby PGN. Approximating the tetrachloric correlation coefficient. *Biometrics* 1983; 39:753–757.
55. *User Menu for Review Manager version 4.2.9: The Cochrane Collaboration, 2006.*
56. Hedges LV, Olkin I. *Statistical methods for meta-analysis.* San Diego, CA: Academic Press, 1985.
57. Cooper H, Hedges LV. *The Handbook of Research Synthesis.* Newbury Park, CA: Russell Sage Foundation, 1994.
58. Evans JD, Bond GR, Meyer PS, Kim HW, Lysaker PH, Gibson PJ, Tunis S. Cognitive and clinical predictors of success in vocational rehabilitation in schizophrenia. *Schizophr Res* 2004; 70:331–342.
59. Gold JM, Goldberg RW, McNary SW, Dixon LB, Lehman AF. Cognitive correlates of job tenure among patients with severe mental illness. *Am J Psychiatry* 2002; 159:1395–1402.
60. Mechanic D, Blider S, McAlpine DD. Employing persons with serious mental illness. *Health Aff (Millwood)* 2002; 21:242–253.
61. Waghorn G, Chant, D. Clinical and non-clinical predictors of vocational recovery for australians with psychotic disorders. *J Rehabil* 2002; 68:40–51.
62. Razzano LA, Cook JA, Burke-Miller JK *et al.* Clinical factors associated with employment among people with severe mental illness: findings from the employment intervention demonstration program. *J Nerv Ment Dis* 2005; 193:705–713.
63. Waghorn G, Chant D, Whiteford H. The strength of self-reported course of illness in predicting vocational recovery for persons with schizophrenia. *J Vocational Rehabil* 2003; 18:33–41.
64. Resnick SG, Rosenheck RA. Recovery and positive psychology: parallel themes and potential synergies. *Psychiatr Serv* 2006; 57:120–122.
65. McGurk SR, Mueser KT. Cognitive and clinical predictors of work outcomes in clients with schizophrenia receiving supported employment services: four-year follow-up. *Adm Policy Ment Health* 2006; 54:1129–1135.
66. Hogarty GE, Flesher S, Ulrich R *et al.* Cognitive enhancement therapy for schizophrenia: effects of a two-year randomized trial on cognition and behavior. *Arch Gen Psychiatry* 2004; 61:866–876.
67. Bond GR, Becker DR, Drake RE *et al.* Implementing supported employment as an evidence-based practice. *Psychiatr Serv* 2001; 52:313–322.
68. Killackey EJ, Jackson HJ, Gleeson J, Hickie IB, McGorry PD. Exciting career opportunity beckons! Early intervention and vocational rehabilitation in first-episode psychosis: employing cautious optimism. *Aust N Z J Psychiatry* 2006; 40:951–996.
69. Lysaker PH, Bond G, Davis LW, Bryson GJ, Bell MD. Enhanced cognitive-behavioral therapy for vocational rehabilitation in schizophrenia: Effects on hope and work. *J Rehabil Res Dev* 2005; 42:673–682.
70. Couture SM, Penn DL, Roberts DL. The functional significance of social cognition in schizophrenia: a review. *Schizophr Bull* 2006; 32:S44–S63.
71. Mueser KT, Salyers MP, Mueser PR. A prospective analysis of work in schizophrenia. *Schizophr Bull* 2001; 27:281–296.
72. Srinivasan L, Tirupati S. Relationship between cognition and work functioning among patients with schizophrenia in an urban area of India. *Psychiatr Serv* 2005; 56:1423–1428.
73. Salkever DS, Karakus MC, Slade EP *et al.* Measures and predictors of community-based employment and earnings of persons with schizophrenia in a multisite study. *Psychiatr Serv* 2007; 58:315–324.
74. Marwaha S, Johnson S, Bebbington P *et al.* Rates and correlates of employment in people with schizophrenia in the UK, France and Germany. *Br J Psychiatry* 2007; 191:30–37.
75. Burke-Miller J.K, Cook JA, Grey D *et al.* Demographic characteristics and employment among people with severe mental illness in a multisite study. *Community Ment Health J* 2006; 42:143–159.
76. Hofer A, Rettenbacher MA, Widschwendter CG, Kemmler G, Hummer M, Fleischhacker WW. Correlates of subjective and functional outcomes in outpatient clinic attendees with schizophrenia

- and schizoaffective disorder. *Eur Arch Psychiatry Clin Neurosci* 2006; 256:246–255.
77. Salkever DS, Slade EP, Karakus MC. Employment retention by persons with schizophrenia employed in non-assisted jobs. *J Rehabil* 2003; 69:19–26.
 78. Latimer EA, Lecomte T, Becker DR *et al.* Generalisability of the individual placement and support model of supported employment: results of a Canadian randomised controlled trial. *Br J Psychiatry* 2006; 189:65–73.
 79. Miettunen J, Lauronen E, Veijola J *et al.* Socio-demographic and clinical predictors of occupational status in schizophrenic psychoses: follow-up within the Northern Finland 1966 birth cohort. *Psychiatry Res* 2007; 150:217–225.
 80. Honkonen T, Stengard E, Virtanen M, Salokangas RK. Employment predictors for discharged schizophrenia patients. *Soc Psychiatry Psychiatr Epidemiol* 2007; 42:372–380.
 81. Resnick SG, Neale MS, Rosenheck RA. Impact of public support payments, intensive psychiatric community care, and program fidelity on employment outcomes for people with severe mental illness. *J Nerv Ment Dis* 2003; 191:139–144.
 82. Catty J, Lissouba P, White S *et al.* Predictors of employment for people with severe mental illness: results of an international six-centre randomised controlled trial. *Br J Psychiatry* 2008; 192:224–231.
 83. Harvey C, Fossey E, Jackson H, Shimitras L. Time use of people with schizophrenia living in North London: predictors of participation in occupations and their implications for improving social inclusion. *J Ment Health* 2006; 15:43–55.
 84. McGurk SR, Mueser KT, Harvey PD, LaPuglia R, Marder J. Cognitive and symptom predictors of work outcomes for clients with schizophrenia in supported employment. *Psychiatr Serv* 2003; 54:1129–1135.